

Memantine improves memory and hippocampal proliferation in adult male rats

Maha Elbeltagy^{1,2}, Duha A. Atieh³, Basil H. Abdin³, Kenan A. S-Yasin³, Ahmad M. Abdulraheem³, Doaa Qattan³, Ahmed S. Salman^{1,2}

¹Department of Anatomy and Histology, Faculty of Medicine, University of Jordan, Amman, Jordan, ²Department of Anatomy, Faculty of Medicine, Menoufia University, Egypt, ³Faculty of Medicine, University of Jordan, Amman, Jordan

Folia Neuropathol 2021; 59 (2): 143-151

DOI: <https://doi.org/10.5114/fn.2021.107607>

Abstract

Neurogenesis occurs during the embryological development of the brain. However, it is universally accepted that in all adult mammalian brains, there are two sites of high-density cell division: the subventricular zone of the lateral ventricles (SVZ) and the subgranular zone (SGZ) of the dentate gyrus of the hippocampal formation.

Doxorubicin (DOX) is an anthracycline agent which results in cognitive deterioration and memory impairment, whereas memantine (MEM) is an NMDA receptor antagonist which is approved for the treatment of Alzheimer's dementia. Many studies have revealed MEM's positive impact on memory and demonstrated that it stimulates neuronal division in the hippocampus.

This study aimed to assess the effect of MEM on spatial memory and neural proliferation in the hippocampus in adult male rats treated with DOX. For this purpose, forty male Sprague-Dawley rats were divided into four groups of ten rats each according to the agent: control, MEM (2.5 mg/kg), DOX (2 mg/kg), and DOX with MEM. The rats were given seven intraperitoneal injections every other day. We tracked the rat's weights to assess the weight-reducing effects of the drugs. In order to test spatial memory, the rats were subjected to the novel location recognition (NLR) task 30 minutes after the last injection. Additionally, Ki67 immunohistochemistry was performed to examine hippocampal proliferation.

The results showed a significant reduction in discrimination index (DI) in the DOX-treated group compared to MEM- ($p < 0.001$) and MEM with DOX-treated groups ($p < 0.001$). There was a significant increase in Ki67-positive cells in the MEM-treated group compared to the saline-treated group. Treatment with DOX impaired hippocampal proliferation compared to treatment with MEM or saline. The co-administration of MEM with DOX ameliorated the decline in hippocampal proliferation compared to treatment with DOX alone. There was a significant weight reduction in the DOX group in comparison to the control group, but MEM attenuated DOX-induced weight loss.

Rats treated with DOX displayed a drop in memory, hippocampal proliferation, and weight compared to the MEM-treated group, whereas the co-administration of MEM with DOX protected memory, hippocampal proliferation, and doxorubicin-induced weight loss.

Key words: memantine, doxorubicin, memory, hippocampus, neurogenesis, novel location recognition.

Communicating author:

Dr Maha Elbeltagy, Faculty of Medicine, University of Jordan, Queen Rania St., Amman, Jordan, phone: +962776336325, e-mail: mahabeltagy76@yahoo.co.uk

Introduction

According to the World Health Organization (WHO), cancer is the second leading cause of death worldwide after cardiovascular diseases, and it is expected to cause 13.1 million deaths in 2030 [51,56]. Doxorubicin (DOX) is a chemotherapeutic agent that is classified as an anthracycline antitumor antibiotic. It is used in the treatment of many cancers including lung cancer, breast cancer, gastric cancer, paediatric cancer, and multiple myeloma [61].

An estimated 17-75% of cancer patients receiving chemotherapy experience cognitive impairment stemming from agents including DOX [29]. Some patients have shown improved cognition over time after the completion of chemotherapy treatment; however, more than 50% continue to experience impairment [53]. In addition, some animal studies using the novel location recognition (NLR) task have found that chronic exposure to DOX impairs memory function consistently with disrupting hippocampal neurogenesis [8].

The hippocampus, a formation of densely packed neurons in the limbic lobe, plays a fundamental role in learning, memory, and spatial navigation [3,46,60]. It facilitates learning *via* its connections to the neocortex; in particular, it is involved in acquiring new memories and then solidifying them, thus transforming short-term memory into long-term memory. The hippocampus itself is composed of two major grey matter elements: the cornu ammonis and the dentate gyrus [22]. Each component consists of distinct types of cells that, together with the entorhinal cortex, interact with each other through circuits and contribute to the learning and memory process [2,19,30,49,50]. Furthermore, high expression of NMDA receptors on pyramidal cells is observed in the hippocampus, which are required for spatial memory [44,45].

It is universally accepted that in all adult mammalian brains, there are two sites of high-density cell division: the subventricular zone of the lateral ventricles (SVZ) and the subgranular zone (SGZ) of the dentate gyrus of the hippocampal formation [18]. Hippocampal neurogenesis in adults is of particular interest in our research since it has been shown to significantly influence cognition and the formation of new memories, thereby considerably affecting learning. Several papers have shown that adult hippocampal neurogenesis improves memory and cognition [15,20,33]. Adult hippocampal neurogenesis has also

been associated with improved pattern separation, which is increasingly recognized as the underlying mechanism of memory and learning [9,12,58]. Furthermore, brains experiencing a reduction in neurogenesis have shown decreased memory performance, whereas those with increased rates of neurogenesis have displayed increases in cognitive function [13,65].

Memantine (MEM) acts as an NMDA receptor antagonist. Previous studies have shown that NMDA receptor antagonists enhance neurogenesis in the brains of adult rats. Accordingly, MEM is being prescribed by clinicians for the treatment of Alzheimer's disease (AD). The excitotoxic process that results from overactive NMDA receptors, which is mediated by the excessive influx of calcium during a sustained release of glutamate, has been associated with strokes, trauma, and chronic degenerative diseases such as AD [11]. Memantine has shown a functional role in the improvement of memory and learning processes after neuronal damage, and it prevents the damage from progressing [5,11].

Memantine acts as a neuroprotective agent rather than a disease-reversing agent [64]. In addition, since it has low affinity to the receptor-associated ion channel, it detaches from its binding site relatively quickly. This results in the alleviation of undesired adverse effects as it is not active for prolonged time periods [35]. Hence, we can deduce that MEM does not alter normal brain signalling and can be predicted to be well tolerated in clinical trials [36]. Due to these properties and its unique mechanism of action, clinical trials have been initiated to study the effects of MEM on other forms of dementia, depression, glaucoma, and severe neuropathic pain [36]. Furthermore, MEM's effect on malignant diseases, like breast and prostate cancer, is being investigated [1,59].

Memantine offers hope for improved quality of life for patients by preventing or at least minimizing the toxic effects of DOX on neural cells and memory.

The current study examined the effects of MEM on both memory and hippocampal proliferation in DOX-treated adult male rats.

Material and methods

Ethics statement

All experiments and animal care were performed in accordance with the University of Jordan's guidelines and with the approval of the local ethics committee.

Animals and drug preparations

Forty male Sprague-Dawley rats (190-225 g) were bought from the University of Jordan's animal office and randomly allocated to four groups: control ($n = 10$), MEM ($n = 10$), DOX ($n = 10$), and DOX with MEM ($n = 10$). The animals were allowed to habituate for two weeks prior to drug administration.

Rats in the MEM group were administered 7 i.p. 2.5 mg/kg doses of memantine (Lundbeck, Denmark) every other day. This dose was modified from a study conducted by Cole *et al.* [10]. Rats in the control group were given an identical volume of 0.9% sterile saline (i.p.). Rats in both the DOX and DOX with MEM groups were administered 7 i.p. 2 mg/kg doses of doxorubicin (EBEWE Pharma, Egypt) every other day. The rats were provided with a 12-hour light/dark cycle (7.00/19.00 h) and free access to food and water.

Behavioural testing

Novel location recognition

In order to test spatial memory, the rats were subjected to the NLR task 30 minutes after the last injection [37]. The NLR task was a spatial variant of a two-trial object recognition task adapted from Dix and Aggleton [14] (see Fig. 1). The apparatus consisted of an arena (a semi-transparent perspex box 49 cm wide \times 66 cm long \times 40 cm high) and objects (pink, weighted water bottles 15 cm high and 7 cm in diameter). The boxes and the water bottles were cleaned with 20% ethanol prior to each experiment and between trials to remove olfactory cues. A square black card was displayed on the wall of the room during the trials to provide prominent cues for spatial orientation.

This apparatus was modified from a previous protocol [14] and was recorded by video camcorder as in our previous study [48]. The procedure consisted of habituating the animals for one hour in the box on the day prior to testing. The following day, as a familiarization trial, two identical objects (water bot-

tlies) were placed in separate locations in the box, and the animals were allowed three minutes to explore. The animals were returned to their home cage for a five-minute inter-trial interval, during which the box was cleaned with 20% ethanol. In the choice trial, the animals were returned to the box for three minutes. One object remained in its original position (the familiar location), while the other object was moved to a new position (the novel location) (see Fig. 1).

The rat was considered to be exploring the object when it sniffed, licked, or chewed the object or directed its nose at a distance ≤ 1 cm from the object [48]. Exploration was scored based on the total time spent on each object (familiar and novel locations). Data were converted to discrimination indices (DI) which means the time spent exploring the novel object minus the time spent exploring the familiar object divided by total exploration time [6,14,16].

Histology and immunohistochemistry

The day after behavioural testing was completed, the rats were put down by rapid stunning and cervical dislocation. Their brains were extracted, trimmed, and fixed in 3% glutaraldehyde overnight. The next day the brains were sectioned using a Leica vibrating microtome. The 4 μ m sections were placed onto positively charged slides for routine staining with haematoxylin and eosin and for Ki67 immunohistochemical analysis. The tissues were dewaxed with xylene and rehydrated through a series of graded ethanols. To retrieve the antigens, the samples were autoclaved in 0.01 M sodium citrate, pH 6.0, at 100°C for 20 minutes and then were heated in a microwave oven (800 W) for 5 minutes [55]. Endogenous peroxidase activity was quenched by incubating the slides in H₂O₂ (3% in methanol) at room temperature for 20 minutes. Non-specific immunoglobulin binding was blocked with 3% bovine serum albumin (manufactured by Merck) in phosphate buffer solution (PBS) at 37°C for 20 minutes.




Fig. 1. Schematic representation of the novel location recognition task protocol.

The polyclonal antibody against Ki67 was bought from Thermofisher, Cat. RB-9043. The primary antibody was diluted in phosphate-buffered saline at a dilution rate of 1 to 50 and incubated at 4°C for 1 h followed by 10 minutes incubation with Goat anti-Rabbit secondary antibody, Thermofisher Cat. A32732 (1 : 250) in PBS for 10 minutes. The slides were counterstained with Mayer's haematoxylin [55].

A systemic random sampling technique [41] was used to choose every twentieth section throughout the length of the dentate gyrus, selecting 10 sections in total. A Zeiss Primo Star microscope (Oberkochen, Germany) equipped with a Canon EOS 550D camera (Tokyo, Japan) was used to confirm the integrity of the selected sections and for counting the proliferating cells.

Counting was done by two independent observers using a double-blind method. Count of Ki67-positive cells was carried out within the SGZ, defined as the zone within three cell diameters of the inner edge of the dentate gyrus (see Fig. 2). Counts from all sections of one dentate gyrus were averaged and multiplied by twenty to provide an estimate of the total number of positive cells in the dentate gyrus [16].

Statistical analysis

Statistical analysis was undertaken and graphs were created using GraphPad Prism 4.0. $P < 0.05$ was regarded as significant. Student's paired *t*-tests were used to

compare the exploration times for rats in each group in the NLR task choice trials. A one-way ANOVA with Bonferroni's post-test was used to compare the number of Ki67-positive proliferating cells and discrimination indices between groups, and a two-way ANOVA with Bonferroni's post-test was used to compare the replicate means of the rat's weights over the injection period between all groups.

Results

The effect of treatment on the novel location recognition task

The NLR task shows interactions with objects either in familiar or novel locations within a test arena. During the familiarization trial, in which the rats explored two identical objects, both the control and the treatment groups showed no preference for either object in terms of the total exploration time (data not shown). During the choice trial, in which one object had been moved to a new location, saline, MEM and MEM with DOX injected groups all explored the novel object significantly more than the old location while rats in the DOX group failed to differentiate between the two locations (data not shown). The discrimination index was calculated as the time spent exploring the novel object minus the time spent exploring the familiar object divided by total exploration time and compared between groups (Fig. 3). There was a significant reduction in discrimination index (DI)




Fig. 2. Representative micrographs of anti-Ki67 immunostaining in the dentate gyrus of **A**) control group, **B**) DOX group, **C**) MEM group, **D**) DOX with MEM group. Ki67-positive cells (arrows) appear dark, indicating proliferation (scale bar = 100 μ m).

in the DOX-treated group compared to MEM- ($p < 0.001$) and MEM with DOX-treated groups ($p < 0.001$).

These findings indicate that the rats treated with MEM along with DOX had protected memory compared to those treated with DOX only and that, compared to MEM treatment, treatment with DOX impaired hippocampal recent memory.

The effect of treatments on proliferating cell counts

There was a significant increase in the total number of Ki67-positive cells in the MEM group compared to the control group ($p < 0.05$). DOX treatment impaired hippocampal proliferation compared to both saline and MEM treatment ($p < 0.001$) for both. The DOX with MEM group showed increased hippocampal proliferation compared to the DOX group ($p < 0.01$). Figure 4 shows these findings correlated with the results obtained from the NLR task.

The effect of different treatments on the rats' weight

As shown in Figure 5, there was significant difference between groups due to both treatment ($p < 0.0001$) and injection periods, which are indicated by arrows ($p < 0.0001$). DOX reduced the weights of the rats after each injection. Overall, there was a significant reduction in weight for rats treated with DOX compared to rats treated with saline throughout the injection period ($p < 0.0001$). Memantine attenuated this weight loss ($p < 0.0001$).

Discussion

This study aimed to assess the effect of MEM on spatial memory and neural proliferation in hippocampus in adult male rats treated with DOX.

Previous animal studies have shown that chronic administration of MEM improved spatial cognitive function as evidenced by decreased errors in a Morris water maze [4,42]. This was also suggested to be true in a transgenic mouse model of AD [43]. Pietá Dias *et al.* demonstrated that chronic MEM administration (20 mg/kg i.p.) over three weeks decreased age-induced spatial memory deficits as investigated through the NLR task. Different test regimens and doses may elucidate this controversy [54].

Enhanced hippocampal proliferation with MEM use was evident in the literature. A study has shown that a single intraperitoneal dose of 50 mg/kg stimu-




Fig. 3. Discrimination index between groups (DI). The doxorubicin (DOX)-treated group did not discriminate between the familiar and novel location. The DI of the group treated with DOX along with memantine (MEM) was improved compared to DOX treatment ($p < 0.001$). The DI of the MEM-treated group was significantly higher than in the DOX-treated group ($p < 0.001$).




Fig. 4. Effect of treatment on the proliferating cell count. There was a significant difference in the total number of Ki67-positive cells between all groups (one-way ANOVA) ($p < 0.0001$, $f = 31.00$).

lated hippocampal dentate gyrus proliferation in both young and elderly rats [38]. A similar effect was found by Jin *et al.* [28].

Notably, in our study, the group of rats given seven i.p. injections of DOX failed to recognize the




Fig. 5. Animal body weight during the injection period between groups. There was a significant reduction in body weight of rats treated with doxorubicin (DOX) compared to rats treated with saline throughout the injection period ($p < 0.0001$). Memantine (MEM) attenuated this weight loss ($p < 0.0001$). Injections are indicated by arrows. The analysis was performed using a two-way ANOVA with Bonferroni's post-test to compare replicate means on a GraphPad Prism 4.

novel location. This finding agrees with previous research showing that DOX disrupted many hippocampal-based memory functions including spatial memory and contextual conditioned fear memory [8,31,63]. In contrast, Fremouw *et al.* found that DOX-treated mice showed normal contextual fear memory and normal performance in the NLR task when compared to the control group [17]. This may be due to the different dosing regimens and differences in neuronal processes and biological factors between rats and mice.

In concordance with our study, DOX-induced cognitive decline may be attributed to decreased hippocampal neurogenesis [8], although one study showed that DOX administration alone did not significantly alter hippocampal neurogenesis unless co-administered with cyclophosphamide [31].

Although the co-administration of MEM with DOX ameliorated DOX-induced cognitive decline and inhibition of hippocampal proliferation, no causal relationship can be deduced based on this observation. Memantine was shown to have other neuroprotective effects like enhancement of synaptic plasticity [67]

and inhibition of apoptosis [27]. In addition, recent *in vitro* studies have demonstrated that MEM ameliorated DOX-induced apoptosis in different types of neuronal cell cultures [25,26]. So further research is encouraged to investigate a causal relationship.

The molecular basis of MEM-induced hippocampal proliferation has not been made clear, although one plausible mechanism is through enhanced BDNF local expression and signalling. This hypothesis is based on a previous finding that MEM caused a dose-related increase in the expression of the BDNF gene, a member of the neurotrophin family, and its receptor TrkB in many cortical regions including the hippocampus [40]. Increasing evidence suggests that increased BDNF levels promote hippocampal neurogenesis in response to different stimuli [24,39,57]. Recent studies have demonstrated that enhanced expression of BDNF contributed to MEM-related enhancement of synaptic plasticity [67] and to the anti-apoptotic effect of MEM [27], but further research is needed to elaborate the effect of BDNF on MEM-induced hippocampal proliferation.

Different NMDA receptor antagonists, including MEM, have shown great promise in reversing chemotherapy-induced cognitive deficits in animal models. A study revealed that methotrexate-induced spatial cognitive impairment, which was attributed in part to increased levels of the excitotoxic glutamate analogue homocysteic acid, was ameliorated by the co-administration of MEM [10]. Another NMDA receptor antagonist, Dextromethorphan, has also been shown to decrease negative cognitive outcomes in methotrexate-treated rats [62]. One study showed that MEM mitigated cisplatin-induced impaired performance of rats in the Morris water maze, and it attenuated the cisplatin-related reduction of the expression of PSD95 and ERK1/2 proteins, which are essential for the formation and maintenance of synaptic plasticity [7]. These preclinical data support the hypothesis that NMDA receptor antagonists like MEM may be used for the treatment of chemotherapy-induced cognitive deficits.

Reduction in body weight is a toxic effect associated with the administration of DOX [21,23,34,66]. We noticed a significant weight reduction in the DOX-injected rats in comparison to the control group during the two-week injection period. However, MEM co-treatment effectively improved DOX-induced weight loss. Accordingly, MEM could be a potential agent for the attenuation and prevention

of weight loss induced by DOX in clinical practice. This can be partly supported by a randomized controlled trial done on Alzheimer's patients showing that MEM was associated with a significant increase in body weight when compared to placebo [52]. In addition, two studies have shown that MEM attenuated weight loss in animal models of Huntington's disease and ulcerative colitis [32,47].

Conclusions

Rats treated with DOX showed a deterioration in memory and an inhibition of cellular proliferation in the hippocampus, in addition to a noticeable decrease in weight. In contrast, the co-administration of MEM and DOX revealed a significant enhancement of memory, a promotion of hippocampal proliferation, plus a remarkable improvement of DOX-induced weight loss.

Acknowledgements

This work is funded by the deanship of scientific research at the University of Jordan, Jordan.

Disclosure

The authors report no conflict of interest.

References

- Albayrak G, Konac E, Dikmen AU, Bilen CY. Memantine induces apoptosis and inhibits cell cycle progression in LNCaP prostate cancer cells. *Hum Exp Toxicol* 2018; 37: 953-958.
- Amaral DG, Witter MP. The three-dimensional organization of the hippocampal formation: a review of anatomical data. *Neuroscience* 1989; 31: 571-591.
- Anand K, Dhikav V. Hippocampus in health and disease: an overview. *Ann Indian Acad Neurol* 2012; 15: 239-246.
- Barnes CA, Danysz W, Parsons CG. Effects of the Uncompetitive NMDA receptor antagonist memantine on hippocampal long-term potentiation, short-term exploratory modulation and spatial memory in awake, freely moving rats. *Eur J Neurosci* 1996; 8: 565-571.
- Beier MT. Treatment strategies for the behavioral symptoms of Alzheimer's disease: Focus on early pharmacologic intervention. *Pharmacotherapy* 2007; 27: 399-411.
- Bruel-Jungerman E, Laroche S, Rampon C. New neurons in the dentate gyrus are involved in the expression of enhanced long-term memory following environmental enrichment. *Eur J Neurosci* 2005; 21: 513-521.
- Cheng J, Liu X, Cao L, Zhang T, Li H, Lin W. Neo-adjuvant chemotherapy with cisplatin induces low expression of NMDA receptors and postoperative cognitive impairment. *Neurosci Lett* 2017; 637: 168-174.
- Christie L-A, Acharya MM, Parihar VK, Nguyen A, Martirosian V, Limoli CL. Impaired cognitive function and hippocampal neurogenesis following cancer chemotherapy. *Clin Cancer Res* 2012; 18: 1954-1965.
- Clelland CD, Choi M, Romberg C, Clemenson GD, Fragniere A, Tyers P, Jessberger S, Saksida LM, Barker RA, Gage FH, Bussey TJ. A functional role for adult hippocampal neurogenesis in spatial pattern separation. *Science* 2009; 325: 210-213.
- Cole PD, Vijayanathan V, Ali NF, Wagshul ME, Tanenbaum EJ, Price J, Dalal V, Gulinello ME. Memantine protects rats treated with intrathecal methotrexate from developing spatial memory deficits. *Clin Cancer Res* 2013; 19: 4446-4454.
- Cosman KM, Boyle LL, Porsteinsson AP. Memantine in the treatment of mild-to-moderate Alzheimer's disease. *Expert Opin Pharmacother* 2007; 8: 203-214.
- Creer DJ, Romberg C, Saksida LM, Van Praag H, Bussey TJ. Running enhances spatial pattern separation in mice. *Proc Natl Acad Sci U S A* 2010; 107: 2367-2372.
- Deng W, Aimone JB, Gage FH. New neurons and new memories: How does adult hippocampal neurogenesis affect learning and memory? *Nat Rev Neurosci* 2010; 11: 339-350.
- Dix SL, Aggleton JP. Extending the spontaneous preference test of recognition: evidence of object-location and object-context recognition. *Behav Brain Res* 1999; 99: 191-200.
- Dupret D, Revest JM, Koehl M, Ichas F, De Giorgi F, Costet P, Abrous DN, Piazza PV. Spatial relational memory requires hippocampal adult neurogenesis. *PLoS One* 2008; 3: e1959.
- ElBeltagy M, Mustafa S, Umka J, Lyons L, Salman A, Gloria Tu CY, Bhalla N, Bennett G, Wigmore PM. Fluoxetine improves the memory deficits caused by the chemotherapy agent 5-fluorouracil. *Behav Brain Res* 2010; 208: 112-117.
- Fremouw T, Fessler CL, Ferguson RJ, Buguete Y. Preserved learning and memory in mice following chemotherapy: 5-Fluorouracil and doxorubicin single agent treatment, doxorubicin-cyclophosphamide combination treatment. *Behav Brain Res* 2012; 226: 154-162.
- Gage FH. Mammalian neural stem cells. *Science* 2000; 287: 1433-1438.
- Gold AE, Kesner RP. The role of the CA3 subregion of the dorsal hippocampus in spatial pattern completion in the rat. *Hippocampus* 2005; 15: 808-814.
- Gould E, Beylin A, Tanapat P, Reeves A, Shors TJ. Learning enhances adult neurogenesis in the hippocampal formation. *Nat Neurosci* 1999; 2: 260-265.
- Hao G, Yu Y, Gu B, Xing Y, Xue M. Protective effects of berberine against doxorubicin-induced cardiotoxicity in rats by inhibiting metabolism of doxorubicin. *Xenobiotica* 2015; 45: 1024-1029.
- Hayman LA, Fuller GN, Cavazos JE, Pflieger MJ, Meyers CA, Jackson EF. The hippocampus: Normal anatomy and pathology. *Am J Roentgenol* 1998; 171: 1139-1146.
- Herman EH, Zhang J, Chadwick DP, Ferrans VJ. Comparison of the protective effects of amifostine and dexrazoxane against the toxicity of doxorubicin in spontaneously hypertensive rats. *Cancer Chemother Pharmacol* 2000; 45: 329-334.
- Hsiao YH, Hung HC, Chen SH, Gean PW. Social interaction rescues memory deficit in an animal model of Alzheimer's disease

- by increasing BDNF-dependent hippocampal neurogenesis. *J Neurosci* 2014; 34: 16207-16219.
25. Jantas D, Lason W. Protective effect of memantine against doxorubicin toxicity in primary neuronal cell cultures: influence a development stage. *Neurotox Res* 2009; 15: 24-37.
 26. Jantas D, Pytel M, Mozrzymas JW, Leskiewicz M, Regulska M, Antkiewicz-Michaluk L, Lason W. The attenuating effect of memantine on staurosporine-, salsolinol- and doxorubicin-induced apoptosis in human neuroblastoma SH-SY5Y cells. *Neurochem Int* 2008; 52: 864-877.
 27. Jantas D, Szymanska M, Budziszewska B, Lason W. An involvement of BDNF and PI3-K/Akt in the anti-apoptotic effect of memantine on staurosporine-evoked cell death in primary cortical neurons. *Apoptosis* 2009; 14: 900-912.
 28. Jin K, Xie L, Mao XO, Greenberg DA. Alzheimer's disease drugs promote neurogenesis. *Brain Res* 2006; 1085: 183-188.
 29. Keeney JTR, Ren X, Warriar G, Noel T, Powell DK, Brelsfoard JM, Sultana R, Saatman KE, St. Clair DK, Butterfield DA. Doxorubicin-induced elevated oxidative stress and neurochemical alterations in brain and cognitive decline: Protection by MESNA and insights into mechanisms of chemotherapy-induced cognitive impairment ("chemobrain"). *Oncotarget* 2018; 9: 30324-30339.
 30. Kesner RP. Behavioral functions of the CA3 subregion of the hippocampus. *Learn Mem* 2007; 14: 771-781.
 31. Kitamura Y, Hattori S, Yoneda S, Watanabe S, Kanemoto E, Sugimoto M, Kawai T, Machida A, Kanzaki H, Miyazaki I, Asanuma M, Sendo T. Doxorubicin and cyclophosphamide treatment produces anxiety-like behavior and spatial cognition impairment in rats: possible involvement of hippocampal neurogenesis via brain-derived neurotrophic factor and cyclin D1 regulation. *Behav Brain Res* 2015; 292: 184-193.
 32. Lee ST, Chu K, Park JE, Kang L, Ko SY, Jung KH, Kim M. Memantine reduces striatal cell death with decreasing calpain level in 3-nitropropionic model of Huntington's disease. *Brain Res* 2006; 1118: 199-207.
 33. Leuner B, Gould E, Shors TJ. Is there a link between adult neurogenesis and learning? *Hippocampus* 2006; 16: 216-224.
 34. de Lima Junior EA, Yamashita AS, Pimentel GD, De Sousa LGO, Santos RVT, Gonçalves CL, Streck EL, de Lira FS, Rosa Neto JC. Doxorubicin caused severe hyperglycaemia and insulin resistance, mediated by inhibition in AMPK signalling in skeletal muscle. *J Cachexia Sarcopenia Muscle* 2016; 7: 615-625.
 35. Lipton S. Pathologically-activated therapeutics for neuroprotection: mechanism of NMDA receptor block by memantine and S-nitrosylation. *Curr Drug Targets* 2007; 8: 621-632.
 36. Lipton S. The molecular basis of memantine action in Alzheimers disease and other neurologic disorders: low-affinity, uncompetitive antagonism. *Curr Alzheimer Res* 2005; 2: 155-165.
 37. Lueptow LM. Novel object recognition test for the investigation of learning and memory in mice. *J Vis Exp* 2017; 2017: 1-9.
 38. Maekawa M, Namba T, Suzuki E, Yuasa S, Kohsaka S, Uchino S. NMDA receptor antagonist memantine promotes cell proliferation and production of mature granule neurons in the adult hippocampus. *Neurosci Res* 2009; 63: 259-266.
 39. Marlatt MW, Potter MC, Lucassen PJ, van Praag H. Running throughout middle-age improves memory function, hippocampal neurogenesis, and BDNF levels in female C57BL/6J mice. *Dev Neurobiol* 2012; 72: 943-952.
 40. Marvanová M, Lakso M, Pirhonen J, Nawa H, Wong G, Castrén E. The neuroprotective agent memantine induces brain-derived neurotrophic factor and trkB receptor expression in rat brain. *Mol Cell Neurosci* 2001; 18: 247-258.
 41. Mayhew TM, Burton GJ. Methodological problems in placental morphometry: apologia for the use of stereology based on sound sampling practice. *Placenta* 1988; 9: 565-581.
 42. Minkeviciene R, Banerjee P, Tanila H. Cognition-enhancing and anxiolytic effects of memantine. *Neuropharmacology* 2008; 54: 1079-1085.
 43. Minkeviciene R, Banerjee P, Tanila H. Memantine improves spatial learning in a transgenic mouse model of Alzheimer's disease. *J Pharmacol Exp Ther* 2004; 311: 677-682.
 44. Morris RGM, Garrud P, Rawlins JNP, O'Keefe J. Place navigation impaired in rats with hippocampal lesions. *Nature* 1982; 297: 681-683.
 45. Morris RGM, Hagan JJ, Rawlins JNP. Allocentric spatial learning by hippocampectomised rats: a further test of the "spatial mapping" and "working memory" theories of hippocampal function. *Q J Exp Psychol Sect B* 1986; 38: 365-395.
 46. Moser EI, Kropff E, Moser MB. Place cells, grid cells, and the brain's spatial representation system. *Annu Rev Neurosci* 2008; 31: 69-89.
 47. Motaghi E, Hajhashemi V, Mahzouni P, Minaian M. The effect of memantine on trinitrobenzene sulfonic acid-induced ulcerative colitis in mice. *Eur J Pharmacol* 2016; 793: 28-34.
 48. Mustafa S, Walker A, Bennett G, Wigmore PM. 5-Fluorouracil chemotherapy affects spatial working memory and newborn neurons in the adult rat hippocampus. *Eur J Neurosci* 2008; 28: 323-330.
 49. Nakazawa K, McHugh TJ, Wilson MA, Tonegawa S. NMDA receptors, place cells and hippocampal spatial memory. *Nat Rev Neurosci* 2004; 5: 361-372.
 50. Nakazawa K, Quirk MC, Chitwood RA, Watanabe M, Yeckel MF, Sun LD, Kato A, Carr CA, Johnston D, Wilson MA, Tonegawa S. Requirement for hippocampal CA3 NMDA receptors in associative memory recall. *Science* 2002; 297: 211-218.
 51. Padma VV. An overview of targeted cancer therapy. *Biomed* 2015; 5: 1-6.
 52. Peskind ER, Potkin SG, Pomara N, Ott BR, Graham SM, Olin JT, McDonald S. Memantine treatment in mild to moderate Alzheimer disease: a 24-week randomized, controlled trial. *Am J Geriatr Psychiatry* 2006; 14: 704-715.
 53. Philpot RM, Ficken M, Johns BE, Engberg ME, Wecker L. Spatial memory deficits in mice induced by chemotherapeutic agents are prevented by acetylcholinesterase inhibitors. *Cancer Chemother Pharmacol* 2019; 84: 579-589.
 54. Pietá Dias C, Martins de Lima MN, Presti-Torres J, Dornelles A, Garcia VA, Siciliani Scalco F, Rewsaat Guimarães M, Constantino L, Budni P, Dal-Pizzol F, Schröder N. Memantine reduces oxidative damage and enhances long-term recognition memory in aged rats. *Neuroscience* 2007; 146: 1719-1725.
 55. Pisamai S, Rungsipipat A, Kunnasut N, Suriyaphol G. Immunohistochemical expression profiles of cell adhesion molecules, matrix metalloproteinases and their tissue inhibitors in central

- and peripheral neoplastic foci of feline mammary carcinoma. *J Comp Pathol* 2017; 157: 150-162.
56. Rivankar S. An overview of doxorubicin formulations in cancer therapy. *J Cancer Res Ther* 2014; 10: 853-858.
 57. Rossi C, Angelucci A, Costantin L, Braschi C, Mazzantini M, Babbini F, Fabbri ME, Tessarollo L, Maffei L, Berardi N, Caleo M. Brain-derived neurotrophic factor (BDNF) is required for the enhancement of hippocampal neurogenesis following environmental enrichment. *Eur J Neurosci* 2006; 24: 1850-1856.
 58. Sahay A, Scobie KN, Hill AS, O'Carroll CM, Kheirbek MA, Burghardt NS, Fenton AA, Dranovsky A, Hen R. Increasing adult hippocampal neurogenesis is sufficient to improve pattern separation. *Nature* 2011; 472: 466-470.
 59. Seifabadi S, Vaseghi G, Javanmard SH, Omid E, Tajadini M, Zarrin B. The cytotoxic effect of memantine and its effect on cytoskeletal proteins expression in metastatic breast cancer cell line. *Iran J Basic Med Sci* 2017; 20: 41-45.
 60. Squire LR. Memory systems of the brain: a brief history and current perspective. *Neurobiol Learn Mem* 2004; 82: 171-177.
 61. Thorn CF, Oshiro C, Marsh S, Hernandez-Boussard T, McLeod H, Klein TE, Altman RB. Doxorubicin pathways: pharmacodynamics and adverse effects. *Pharmacogenet Genomics* 2011; 21: 440-446.
 62. Vijayanathan V, Gulinello M, Ali N, Cole PD. Persistent cognitive deficits, induced by intrathecal methotrexate, are associated with elevated CSF concentrations of excitotoxic glutamate analogs and can be reversed by an NMDA antagonist. *Behav Brain Res* 2011; 225: 491-497.
 63. Wagawade JD, Roy S, Hullatti K, Khode NR, Pednekar H, Hegde H V, Kholkute SD. Doxorubicin induced cognition impairment in rat model. *Asian J Pharm Clin Res* 2015; 8: 301-304.
 64. Williams BS, Buvanendran A. Memantine. In: Sinatra RS, Jahr JS, Watkins-Pitchford JM, (Eds.). *The Essence of Analgesia and Analgesics*. Cambridge University Press, Cambridge 2010; pp. 319-320.
 65. Zhao C, Deng W, Gage FH. Mechanisms and functional implications of adult neurogenesis. *Cell* 2008; 132: 645-660.
 66. Zhao X, Zhang J, Tong N, Liao X, Wang E, Li Z, Luo Y, Zuo H. Berberine attenuates doxorubicin-induced cardiotoxicity in mice. *J Int Med Res* 2011; 39: 1720-1727.
 67. Zhu G, Li J, He L, Wang X, Hong X. MPTP-induced changes in hippocampal synaptic plasticity and memory are prevented by memantine through the BDNF-TrkB pathway. *Br J Pharmacol* 2015; 172: 2354-2368.